Final Expense Term Whole Life IUL (Fixed) VL (Securities)		
LIFE CHECKUP FORM		
NAME NAME		
Owner: Spouse:		
ADDRESS:		
CITY:ST:ST:SIP:		
PHONE:COUNTY:		
EMAIL:		
Owner DOB #:Spouse DOB #:		
Owner SSN #:Spouse SSN #:		
CURRENT Life Ins. PLANS: Term Whole IUL (Fixed) VL (Securites)		
Replacing a policy: Y/N if YES, name of company, amt. of benefit, policy #, phone # of co		
Prescription Drugs (dosage & how often)/		
Medical Conditions: Smoker/Non-Smoker:		
Diabetes (if insulin are you taking 50 units or more/onset before 9 yrs/insulin before 40/45): Cancer (how long)		
Heart Conditions (COPD/Angina etc):		
Own Home:How much is Owed:		
Mortgage Insurance (explain how it works)		
CHILDREN: How Many Ages:		
DOB: (C1) DOB: (C2) DOB: (C3)		

Do you have a college fund if something were to happen to you?

Underwriting Knockout Medical Conditions for Each Carrier

AIG: Impairments that are automatically declined – do not routinely order an APS

Abdominal Aortic Aneurysm surgically corrected in the last 6 months	Kidney Dialysis
Alcohol treatment in the last 2 years	Mental Disorder (including anxiety or depression requiring hospitalization or disability in last year
Automatic Defibrillator (ACID) implanted with history of Cardiac Arrest	Medical testing advised but not yet completed
Alzheimer's disease or Dementia	MI/heart attack in last 6 months
Blood Pressure exceeding 185/100	Myelodysplastic Syndrome
Cancer treatment (current)	Organ Transplant (awaiting or recipient) (except kidney-contact your underwriter)
Cirrhosis of Liver	Pregnant (current) with Gestational Diabetes, toxemia, eclampsia, pre-eclampsia
COPD/Emphysema, severe (on oxygen or disabling)	Quadriplegia
CVA (stroke)within 1 year	Suicide attempt in the last year
Diabetes if significantly complicated (i.e.	Suicide attempts (more than one) if the last one
amputation etc.) or very poor control	was within the last 2 years
Drug use (other than marijuana) in the last 3 years	Surgery (major) advised but not yet completed
Gastric/Intestinal Bypass within 6 months	Term Insurance, age > 70 & rated over Table D or with a flat extra
Grand Mal Seizures within 1 year of diagnosis	Valve replacement within 6 months
HIV Positive	
Juvenile Onset Diabetes, younger than age 20	

Transamerica: <u>SUBJECT TO UNDERWRITING AND CHANGE WITHOUT NOTICE.</u>

MEDICAL CONDITION	DECISION
AIDS/HIV/ARC	Decline
Alcoholism/Alcohol Abuse – Used or been diagnosed	Within 2 years – Graded
with, treated, tested positive for, or been given medical	Within 2–4 years – Standard
advice by a member of medical profession.	Recovery (no drinks) > 4 years – Preferred
ALS (Lou Gehrig's disease)	Decline
Alzheimer's/Dementia/Memory Loss/Cognitive	Decline
Disorders	
Amputation (excluding at time of accident/trauma)	Decline
Anemia (other than Iron Deficiency) If Sickle Cell	Within last 2 years – Standard
Anemia, please see the section for Sickle Cell Anemia	
Aneurysm	Present (not surgically corrected) or surgically corrected
	within 1 year - Graded
	Surgically corrected between 1–2 years – Standard
	Surgically corrected > 2 years - Preferred
Angina	Onset within 1 year - Graded
	Onset 1–2 years – Standard
	Onset > 2 years – Preferred

Angioplasty (of any kind)	Onset prior age 45 – Graded
Tangle Grand Carry Carry	Onset age 45 and above: Within 1 year – Graded
	Within 2 years – Standard
Arrhythmia	Treatment within 2 years – Standard
•	Condition resolved, no current treatment, last
	treatment > 2 years – Preferred
Assisted Living/Long Term Care Facility – Home	Currently – Decline
healthcare is defined as: medical care provided by a	Within 2 years – Graded
medical professional, friends or family member	
including, but not limited to, arranging medications,	
taking blood pressure or sugar readings, administering	
medications, wound care, feeding tube, etc.	
Asthma (Chronic) – Defined as: Using inhalers year	Standard
round on a daily or weekly basis, or filling inhalers 6 or	
more times in any 12 month period.	
Atrial Fibrillation	Treatment within 2 years – Standard
	Condition resolved, no current treatment, last
	treatment > 2 years – Preferred
Attempted Suicide	Within 2 years – Decline
Autism	Current age 0-17 - Decline
	Mild (Highly Functional): Preferred
	All others – Decline
Bipolar	Current age 0–17 – Decline
	Current age 18–85 – Standard
Black Lung	Standard
Blood Clots (no complications/time since resolved)	Diagnosed and treated within 2 years – Standard
	Diagnosed and treated > 2 years – Preferred
Bone Marrow Transplant (Including donor stem cells)	Decline
Blood Disorder - Anemia (excluding Iron Deficiency	Diagnosed and treated within 2 years – Standard
Anemia/Sickle Cell Anemia), Polycythemia,	No current treatment, last treatment > 2 years – Preferred
Thrombocytopenia, Hemophilia, and other coagulation disorders.	Preferred
Bone Marrow Transplant (Including donor stem cells)	Decline
Bronchitis (Chronic)	Standard
Build Note: If medical questions are Standard and build	See chart pg. 23
is Standard, then you should quote Graded	See chart pg. 23
Bypass	Onset prior age 45 – Graded
Буризэ	Onset age 45 and above: Within 1 year – Graded
	Within 2 years – Standard
	> 2 years – Preferred
Cancer (other than Basal Cell)	Any onset (other than Basal Cell): Within 2 years –
,	Decline
	Metastatic - Decline
	Recurrent - Decline
	Multiple cancers - Decline
	Including lymph node involvement – Decline
	Within last 4 years – Graded
	Prior age 45 (other than Basal Cell) – Standard
Cardiac Surgery	Onset prior age 45 – Graded
	Onset age 45 and above: Within 1 year – Graded
	Within 2 years – Standard
	> 2 years – Preferred

Cardiomyopathy	Standard
Cerebral Palsy	Decline
Chest Pain	See Angina
Chronic Pain – Defined as: Disabled due to pain, using	Currently unemployed/disabled due to pain – Graded
narcotic pain meds on a daily or weekly basis, or 6 fills	Diagnosed with/treated for within 1 year – Graded
of narcotic pain meds in the last 12 month period.	Employed, no disability due to pain or treatment > 1
·	year – Preferred
Circulatory Disorder – (Peripheral Vascular Disease	Onset of PVD, PAD, or CAD prior to age 45 – Graded
(PVD) or Peripheral Arterial Disease (PAD), Phlebitis,	Diagnosed and treated within 2 years – Standard No
Arteriosclerosis, Coronary Artery Disease (CAD))	current treatment, last treatment > 2 years – Preferred
Cirrhosis	Graded
Clotting Disorder	See Blood Clots
Congestive Heart Failure/Heart Failure/Diastolic Heart	Onset prior age 45 – Decline
Failure	Onset age 45–80 – Graded
COPD (Chronic Obstructive Pulmonary Disease)	Standard
Coronary Artery Disease	See Circulatory Disorder
COVID	≤ age 70 Postpone minimum 3 months from date of
	recovery
	> age 70 Postpone minimum 6 months from date of
	recovery
Crohn's Disease	Onset prior to age 26 – Graded
	Onset age > 26 – Standard
Defibrillator Implant	See Pacemaker Implant
Dementia	Decline
Depression	Current age 0–17 – Decline
	Current age 18 and older – Preferred
Dishetes	Onest union and 20 (athough an Contational Bishatas)
Diabetes	Onset prior age 20 (other than Gestational Diabetes) –
Diapetes	Graded
Diapetes	Graded Onset > age 20 with insulin use within 2 years –
Diapetes	Graded Onset > age 20 with insulin use within 2 years — Standard Onset > age 20 oral or diet within 2 years —
Diapetes	Graded Onset > age 20 with insulin use within 2 years –
Diabetes Diabetic Coma	Graded Onset > age 20 with insulin use within 2 years – Standard Onset > age 20 oral or diet within 2 years – Preferred Decline
	Graded Onset > age 20 with insulin use within 2 years — Standard Onset > age 20 oral or diet within 2 years — Preferred Decline Received within 1 year — Graded
Diabetic Coma Dialysis	Graded Onset > age 20 with insulin use within 2 years — Standard Onset > age 20 oral or diet within 2 years — Preferred Decline Received within 1 year — Graded Received within 4 years - Standard
Diabetic Coma Dialysis Diastolic Heart Failure	Graded Onset > age 20 with insulin use within 2 years – Standard Onset > age 20 oral or diet within 2 years – Preferred Decline Received within 1 year – Graded Received within 4 years - Standard See Congestive Heart Failure
Diabetic Coma Dialysis	Graded Onset > age 20 with insulin use within 2 years — Standard Onset > age 20 oral or diet within 2 years — Preferred Decline Received within 1 year — Graded Received within 4 years - Standard See Congestive Heart Failure Donor stem cells — Decline
Diabetic Coma Dialysis Diastolic Heart Failure Donor Stem-Cell Transplant	Graded Onset > age 20 with insulin use within 2 years — Standard Onset > age 20 oral or diet within 2 years — Preferred Decline Received within 1 year — Graded Received within 4 years - Standard See Congestive Heart Failure Donor stem cells — Decline Self-donated stem cells — Preferred
Diabetic Coma Dialysis Diastolic Heart Failure Donor Stem-Cell Transplant Down's Syndrome	Graded Onset > age 20 with insulin use within 2 years – Standard Onset > age 20 oral or diet within 2 years – Preferred Decline Received within 1 year – Graded Received within 4 years - Standard See Congestive Heart Failure Donor stem cells – Decline Self-donated stem cells – Preferred Decline
Diabetic Coma Dialysis Diastolic Heart Failure Donor Stem-Cell Transplant Down's Syndrome Driving (Including: reckless driving, OWI/DUI, or three or	Graded Onset > age 20 with insulin use within 2 years – Standard Onset > age 20 oral or diet within 2 years – Preferred Decline Received within 1 year – Graded Received within 4 years - Standard See Congestive Heart Failure Donor stem cells – Decline Self-donated stem cells – Preferred Decline Within 2 years – Decline
Diabetic Coma Dialysis Diastolic Heart Failure Donor Stem-Cell Transplant Down's Syndrome	Graded Onset > age 20 with insulin use within 2 years — Standard Onset > age 20 oral or diet within 2 years — Preferred Decline Received within 1 year — Graded Received within 4 years - Standard See Congestive Heart Failure Donor stem cells — Decline Self-donated stem cells — Preferred Decline Within 2 years — Decline Within 2 years — Standard
Diabetic Coma Dialysis Diastolic Heart Failure Donor Stem-Cell Transplant Down's Syndrome Driving (Including: reckless driving, OWI/DUI, or three or more moving violations)	Graded Onset > age 20 with insulin use within 2 years – Standard Onset > age 20 oral or diet within 2 years – Preferred Decline Received within 1 year – Graded Received within 4 years - Standard See Congestive Heart Failure Donor stem cells – Decline Self-donated stem cells – Preferred Decline Within 2 years – Decline Within 2-4 years – Standard > 4 years – Preferred
Diabetic Coma Dialysis Diastolic Heart Failure Donor Stem-Cell Transplant Down's Syndrome Driving (Including: reckless driving, OWI/DUI, or three or	Graded Onset > age 20 with insulin use within 2 years – Standard Onset > age 20 oral or diet within 2 years – Preferred Decline Received within 1 year – Graded Received within 4 years - Standard See Congestive Heart Failure Donor stem cells – Decline Self-donated stem cells – Preferred Decline Within 2 years – Decline Within 2-4 years – Standard > 4 years – Preferred Used or received treatment for within 2 years – Graded
Diabetic Coma Dialysis Diastolic Heart Failure Donor Stem-Cell Transplant Down's Syndrome Driving (Including: reckless driving, OWI/DUI, or three or more moving violations)	Graded Onset > age 20 with insulin use within 2 years — Standard Onset > age 20 oral or diet within 2 years — Preferred Decline Received within 1 year — Graded Received within 4 years - Standard See Congestive Heart Failure Donor stem cells — Decline Self-donated stem cells — Preferred Decline Within 2 years — Decline Within 2 years — Standard > 4 years — Preferred Used or received treatment for within 2 years — Graded Used or received treatment for within 4 years —
Diabetic Coma Dialysis Diastolic Heart Failure Donor Stem-Cell Transplant Down's Syndrome Driving (Including: reckless driving, OWI/DUI, or three or more moving violations) Drug Use/Abuse (including prescription drugs)	Graded Onset > age 20 with insulin use within 2 years – Standard Onset > age 20 oral or diet within 2 years – Preferred Decline Received within 1 year – Graded Received within 4 years - Standard See Congestive Heart Failure Donor stem cells – Decline Self-donated stem cells – Preferred Decline Within 2 years – Decline Within 2 years – Standard > 4 years – Preferred Used or received treatment for within 2 years – Graded Used or received treatment for within 4 years – Standard
Diabetic Coma Dialysis Diastolic Heart Failure Donor Stem-Cell Transplant Down's Syndrome Driving (Including: reckless driving, OWI/DUI, or three or more moving violations) Drug Use/Abuse (including prescription drugs)	Graded Onset > age 20 with insulin use within 2 years – Standard Onset > age 20 oral or diet within 2 years – Preferred Decline Received within 1 year – Graded Received within 4 years - Standard See Congestive Heart Failure Donor stem cells – Decline Self-donated stem cells – Preferred Decline Within 2 years – Decline Within 2-4 years – Standard > 4 years – Preferred Used or received treatment for within 2 years – Graded Used or received treatment for within 4 years – Standard See Wheelchair
Diabetic Coma Dialysis Diastolic Heart Failure Donor Stem-Cell Transplant Down's Syndrome Driving (Including: reckless driving, OWI/DUI, or three or more moving violations) Drug Use/Abuse (including prescription drugs) Electric Cart Electric Scooter	Graded Onset > age 20 with insulin use within 2 years — Standard Onset > age 20 oral or diet within 2 years — Preferred Decline Received within 1 year — Graded Received within 4 years - Standard See Congestive Heart Failure Donor stem cells — Decline Self-donated stem cells — Preferred Decline Within 2 years — Decline Within 2 years — Standard > 4 years — Preferred Used or received treatment for within 2 years — Graded Used or received treatment for within 4 years — Standard See Wheelchair
Diabetic Coma Dialysis Diastolic Heart Failure Donor Stem-Cell Transplant Down's Syndrome Driving (Including: reckless driving, OWI/DUI, or three or more moving violations) Drug Use/Abuse (including prescription drugs) Electric Cart Electric Scooter Emphysema	Graded Onset > age 20 with insulin use within 2 years — Standard Onset > age 20 oral or diet within 2 years — Preferred Decline Received within 1 year — Graded Received within 4 years - Standard See Congestive Heart Failure Donor stem cells — Decline Self-donated stem cells — Preferred Decline Within 2 years — Decline Within 2 years — Standard > 4 years — Preferred Used or received treatment for within 2 years — Graded Used or received treatment for within 4 years — Standard See Wheelchair See Wheelchair
Diabetic Coma Dialysis Diastolic Heart Failure Donor Stem-Cell Transplant Down's Syndrome Driving (Including: reckless driving, OWI/DUI, or three or more moving violations) Drug Use/Abuse (including prescription drugs) Electric Cart Electric Scooter Emphysema Encephalitis	Graded Onset > age 20 with insulin use within 2 years — Standard Onset > age 20 oral or diet within 2 years — Preferred Decline Received within 1 year — Graded Received within 4 years - Standard See Congestive Heart Failure Donor stem cells — Decline Self-donated stem cells — Preferred Decline Within 2 years — Decline Within 2-4 years — Standard > 4 years — Preferred Used or received treatment for within 2 years — Graded Used or received treatment for within 4 years — Standard See Wheelchair See Wheelchair Standard Present/treatment within 1 year — Graded
Diabetic Coma Dialysis Diastolic Heart Failure Donor Stem-Cell Transplant Down's Syndrome Driving (Including: reckless driving, OWI/DUI, or three or more moving violations) Drug Use/Abuse (including prescription drugs) Electric Cart Electric Scooter Emphysema Encephalitis Epilepsy	Graded Onset > age 20 with insulin use within 2 years — Standard Onset > age 20 oral or diet within 2 years — Preferred Decline Received within 1 year — Graded Received within 4 years - Standard See Congestive Heart Failure Donor stem cells — Decline Self-donated stem cells — Preferred Decline Within 2 years — Decline Within 2-4 years — Standard > 4 years — Preferred Used or received treatment for within 2 years — Graded Used or received treatment for within 4 years — Standard See Wheelchair See Wheelchair Standard Present/treatment within 1 year — Graded See Seizures
Diabetic Coma Dialysis Diastolic Heart Failure Donor Stem-Cell Transplant Down's Syndrome Driving (Including: reckless driving, OWI/DUI, or three or more moving violations) Drug Use/Abuse (including prescription drugs) Electric Cart Electric Scooter Emphysema Encephalitis	Graded Onset > age 20 with insulin use within 2 years — Standard Onset > age 20 oral or diet within 2 years — Preferred Decline Received within 1 year — Graded Received within 4 years - Standard See Congestive Heart Failure Donor stem cells — Decline Self-donated stem cells — Preferred Decline Within 2 years — Decline Within 2—4 years — Standard > 4 years — Preferred Used or received treatment for within 2 years — Graded Used or received treatment for within 4 years — Standard See Wheelchair Standard Present/treatment within 1 year — Graded See Seizures Awaiting trial — Decline
Diabetic Coma Dialysis Diastolic Heart Failure Donor Stem-Cell Transplant Down's Syndrome Driving (Including: reckless driving, OWI/DUI, or three or more moving violations) Drug Use/Abuse (including prescription drugs) Electric Cart Electric Scooter Emphysema Encephalitis Epilepsy	Graded Onset > age 20 with insulin use within 2 years — Standard Onset > age 20 oral or diet within 2 years — Preferred Decline Received within 1 year — Graded Received within 4 years - Standard See Congestive Heart Failure Donor stem cells — Decline Self-donated stem cells — Preferred Decline Within 2 years — Decline Within 2-4 years — Standard > 4 years — Preferred Used or received treatment for within 2 years — Graded Used or received treatment for within 4 years — Standard See Wheelchair See Wheelchair Standard Present/treatment within 1 year — Graded See Seizures

Heart Attack	Onset prior age 45 – Graded
100107100001	Onset age 45 and above: Within 1 year – Graded
	Within 2 years – Standard > 2 years – Preferred
Heart Failure	See Congestive Heart Failure
Heart Murmur	See Atrial Fibrillation
Heart Valve Replacement	Onset prior age 45 – Graded
Treate valve replacement	Onset age 45 and above: Within 1 year – Graded
	Within 2 years – Standard
	> 2 years – Preferred
Hepatitis A	See Liver Disorder
Hepatitis B	See Liver Disorder
Hepatitis C If Hepatitis C is considered Standard, and	Currently have or been treated within 24 months –
another question in Section 15 is answered yes, then	Graded
you should quote Graded.	Cured, treatment over 24 months – Standard
Hodgkin's/Non-Hodgkin's/Lymphoma	See Cancer
Home Health Care	See Assisted Living
Hospice	Decline Decline
Hospitalization (currently)	See Assisted Living
Huntington's Disease	Decline
Illegal Drugs	Used or received treatment for within 2 years – Graded Used or received treatment for within 4 years –
	Standard
Incarceration	
	Within 2 years – Decline
Iron Deficiency Anemia	Preferred
Irregular Heartbeat	See Arrhythmia
Kidney Disease/Disorder (Chronic Kidney Disease)	Any treatment within 4 years or Stages 1–3 – Standard
= ·1	Stages 4–5 – Graded
Kidney Failure	See Renal Failure
Liver Disease (excluding Fatty Liver Disease)	Diagnosed and/or treatment within 1 year – Graded
	Diagnosed and/or treatment within 2 years – Standard
	Treated and resolved > 2 years – Preferred
Long Term Care	See Assisted Living
Lou Gehrig Disease	See ALS
Lupus	See Systemic Lupus Erythematosus (SLE)
Marijuana	Used within 2 years – Graded
	Used within 4 years – Standard
Memory Loss	See Alzheimer's
Mental Health	See specific diagnosis
Mental Incapacity	Decline
Mental Retardation	Decline
Multiple Sclerosis	Standard
Muscular Dystrophy	Graded
Myocardial Infarction	Onset prior age 45 – Graded
	Onset over age 45: Within 1 year – Graded
	Within 2 years – Standard
	> 2 years: Preferred
Nursing Home	See Assisted Living
Organ Transplant (including donor stem-cell)	Decline
Oxygen	Within 1 yr - Graded.
	Last used > 1 yr Preferred.

Pacemaker/Defibrillator Implant	Prior to age 45 – Graded
Tatemaner, Jenermaner implant	Within 1 year – Graded
	Within 1–2 years and still present (regardless of date
	installed) – Standard
Pancreatitis (nonalcoholic)	Diagnosed and/or treatment within 2 years – Standard
·	Treated and resolved > 2 years – Preferred
Parkinson's Disease	Onset prior age 45 – Graded
	Onset 45–85 – Standard
Parole/Probation (currently)	Within 2 years – Decline
Peripheral Artery Disease (PAD)	Prior age 45 – Graded
Peripheral Vascular Disease (PVD)	Prior age 45 – Graded
Phlebitis	See Blood Clot
Post-Traumatic Stress Disorder (PTSD)	See Depression
Prison (within 2 years)	See Incarceration
Pulmonary Fibrosis	Decline
Pulmonary Hypertension	See Circulatory Disorder
Renal Failure	Stages 4–5 – Graded
Respiratory Disease (Chronic)	See Asthma
Rheumatoid Arthritis	See Chronic Pain
Sarcoidosis	Standard
Schizophrenia	Current age 0–17 – Decline
36m20pmcmu	Current age 18–85 – Standard
Seizures	6+ episodes/attacks within 1 year – Graded 6+
SCIZUICS	episodes/attacks within 2 years – Standard
Sickle Cell Anemia	Decline
Sleep Apnea	CPAP/treatment with supplemental oxygen/oxygen
Sieep Aprilea	concentrator use – Graded
	CPAP/treatment without supplemental oxygen –
	Preferred
Spina Bifida	Present and surgically corrected within 2 years –
Spina Sinaa	Standard Surgically corrected > 2 years – Preferred
Stent Implant	Onset prior age 45 – Graded
	Onset age 45 and above: Within 1 year – Graded
	Within 2 years – Standard
	> 2 years – Preferred
Stroke (single attack/no complications or residuals)	Onset prior age 45 – Graded
of one (single actually no complications of residuals)	Onset age 45 and above: Within 1 year – Graded
	Within 2 years – Standard
	> 2 years – Preferred
Surgery (Been advised or planning to have surgery	Decline
requiring general anesthesia)	2555
Systemic Lupus Erythematosus (SLE)	Diagnosed with or been treated within 2 years – Graded
, , , , , , , , , , , , , , , , , , , ,	Remission, no treatment within 2 years – Standard
Terminal Illness (death expected within 18 months)	Decline Decline
Tissue Transplant	Donor stem cells – Decline
· · · · · · · · ·	Self-donated tissue transplant – Preferred
Transient Ischemic Attack/TIA	Onset prior age 45 – Graded
	Onset age 45 and above: Within 1 year – Graded
	Within 2 years – Standard
	> 2 years – Preferred
Tuberculosis	Current – Standard
	Treated and Resolved > 2 years – Preferred
L	,

Ulcerative Colitis	Standard
Vascular Disorder	Diagnosed and treated within 2 years – Standard
	No current treatment, last treatment > 2 years -
	Preferred
Wheelchair/Electric Scooter/Electric Cart**	Any use within 2 years (current use or use occasionally at facilities such as, but not limited to, the grocery store, department stores, warehouse stores, airports) – Graded Reason for use is expected to resolve in next 3 months or the reason for use has resolved and they are no longer using a wheelchair/ electric scooter/electric cart any longer. – Preferred

Mutual Of Omaha:

Is the proposed insured Currently: Bedridden or confined to any hospital, nursing home, long-term care facility or skilled nursing facility; or receiving or been advised to receive care in a nursing home, hospice care, or home health care?. Requiring assistance with activities of daily living such as taking medications, bathing, dressing, eating, toileting, getting in and out of a chair or bed, or control of bowel or bladder problems? Requiring any of the following (other than for	Has the proposed insured ever been: Diagnosed as having Acquired Immune Deficiency Syndrome (AIDS), AIDS Related Complex (ARC), or Human Immunodeficiency Virus (HIV) Infection (symptomatic or asymptomatic) or been treated for AIDS, ARC, or HIV by a physician or health care provider? Diagnosed with, been treated for or advised by a physician or health care provider to receive treatment for Alzheimer's Disease, Dementia, Huntington's Disease, Sickle Cell Anemia, Myelodysplastic Syndrome (MDS), Lou Gehrig's Disease (ALS), Quadriplegia, Paraplegia, Down's Syndrome, mental incapacity, congestive heart failure, Cirrhosis, Metastatic Cancer or recurrent Cancer of the same type? Diagnosed with insulin shock, diabetic coma, or had
fractures, bone or joint surgery, including replacement): wheelchair, electric scooter, or oxygen equipment to assist breathing (excluding use for sleep apnea)?	an amputation due to diabetic complications or diagnosed with End Stage Renal Disease or requiring dialysis?.
In the past 12 months, has the Proposed Insured been: Advised by a physician to have a surgical operation, diagnostic testing other than for routine screening purposes or for those related to HIV/AIDS, treatment, hospitalization, or other procedure which has not been done or for which results are not known?	Advised to receive or have received an organ or bone marrow transplant?.
Diagnosed by a physician or health care provider as having heart disease or heart surgery of any kind? .	Diagnosed by a physician or health care provider as having a terminal medical condition that is expected to result in death within the next twelve 12 months?.
In the past 2 years, has the Proposed Insured been diagnosed with, been treated for or advised by a physician or health care provider to receive treatment	Has the proposed insured ever been: Diagnosed as having Acquired Immune Deficiency Syndrome (AIDS), AIDS Related Complex (ARC),

for any form of cancer (except basal or squamous cell skin cancer)?	or Human Immunodeficiency Virus (HIV) Infection (symptomatic or asymptomatic) or been treated for AIDS, ARC, or HIV by a physician or health care provider?
Has the Proposed Insured ever (a) received care or treatment for, or (b) been advised by a physician or health care provider to seek treatment for: Diabetes before age 50 or diabetes at any age with complications of Retinopathy (eye), Nephropathy (kidney), Neuropathy (nerve) or Peripheral Vascular Disease (PVD or PAD)?	In the past 2 years, has the Proposed Insured: Been convicted of or currently awaiting trial for a felony?
Hepatitis C?	Been treated for or advised to have treatment for alcohol or drug abuse or convicted more than once of reckless driving or driving under the influence of drugs or alcohol?.
Chronic Lung Disease, including Chronic Obstructive Pulmonary Disease (COPD), Chronic Bronchitis, Emphysema, or Sarcoidosis?	Used unlawful drugs in any form or abused or misused prescription drugs? .
In the past 4 years, has the Proposed Insured: (a) received care or treatment for, or (b) been advised by a physician or health care provider to seek treatment for:	In the past 2 years, has the Proposed Insured been hospitalized by a physician or health care provider for any mental or nervous disorder?
Cancer, Leukemia, Melanoma or any other internal cancer (except basal or squamous cell skin cancer)?	In the past 12 months, has the Proposed Insured consulted a physician for chronic cough, unexplained weight loss greater than 10 pounds, fatigue or unexplained gastrointestinal bleeding?.
Chronic Kidney Disease, Systemic Lupus or Scleroderma?	
Bipolar Depression, Schizophrenia, Parkinson's Disease or Multiple Sclerosis?	
In the past 2 years, has the Proposed Insured: (a) received care or treatment for, or (b) been advised by a physician or health care provider to seek treatment for: Coronary Artery Disease, Heart Attack, Coronary Artery Bypass Surgery, Angioplasty, Cardiomyopathy, irregular heart rhythm, or Valvular Heart Disease with surgical repair or replacement?	
Stroke or Transient Ischemic Attack (TIA)?	